

Angling Indiana - 2009 FISH CONSUMPTION ADVISORY

Kosciusko County- advice for sensitive populations*

Waterbody	Fish Species	Fish Length (inches)	Maximum Amount for Adults to Eat** (Meals)
All Indiana Rivers and Streams (unless specified)	Carp	ALL	0
Center Lake	Black Bullhead	11+	0
	Bluegill	7+	0
	Largemouth Bass	14+	0
Dewart Lake	Black Crappie	Up to 12	8 ounces per week (1 meal/week)
	Bluegill	Up to 8	8 ounces per week (1 meal/week)
	Northern Pike	30+	0
Easterday Ditch	Carp	Up to 23	2 ounces per week (1 meal/month)
		23+	0
Lake Wawasee	Bullhead	15+	0
Little Barbee Lake	Bluegill	Up to 7	8 ounces per week (1 meal/week)
Palestine Lake	Bluegill	8+	0
	Largemouth Bass	12+	0
Pike Lake	Largemouth Bass	11+	0
	Walleye	14+	0
Tippecanoe Lake	Largemouth Bass	12+	0
Tippecanoe River (downstream of State Road 15)	Bluegill	6+	0
	Carp	20+	0
	Redhorse Species	16+	0
Tippecanoe River (Oswego to State Road 15)	Bluegill	Up to 5	8 ounces per week (1 meal/week)
	Carp	Up to 23	2 ounces per week (1 meal/month)
		23+	0
	Longear Sunfish	Up to 5	8 ounces per week (1 meal/week)
	Rock Bass	Up to 6	8 ounces per week (1 meal/week)
Warmouth	Up to 6	8 ounces per week (1 meal/week)	
Winona Lake	Bluegill	Up to 8	8 ounces per week (1 meal/week)
	Carp	24+	0
	Largemouth Bass	12+	0
	Walleye	24+	0
	White Bass	15+	0
	White Sucker	19+	0
Yellow Perch	Up to 8	8 ounces per week (1 meal/week)	

] *Sensitive populations include: pregnant or nursing women, women that will become pregnant, and children under 6 years of age. These consumers should use caution when eating some types of sportfish.

**The maximum amount for adults to eat is determined by fishing location, species and fish length. For instance, if you eat the maximum amount of one species during a week, you should not eat any other fish which has a consumption limitation until the following week. Example - if you eat the limit (4 ounces) of a 20 inch long Carp from the Easy Catch River, then you should not eat any other sportfish from the list until the following week.